

Black holes on cylinders are not algebraically special

Pieter-Jan De Smet*

*Instituut voor theoretische fysica, Katholieke Universiteit Leuven,
Celestijnenlaan 200D, B-3001 Leuven, Belgium.*
E-mail: Pieter-Jan.DeSmet@fys.kuleuven.ac.be

ABSTRACT: We give a Petrov classification for five-dimensional metrics. We classify Ricci-flat metrics that are static, have an $SO(3)$ isometry group and have Petrov type 22. We use this classification to look for the metric of a black hole on a cylinder, i.e. a black hole with asymptotic geometry four-dimensional Minkowski space times a circle. Although a black string wrapped around the S^1 and the five-dimensional black hole are both algebraically special, it turns out that the black hole on a cylinder is not.

*Address after 1 September 2002: SUNY at Stony Brook, USA

Contents

1. Introduction	1
2. Petrov classification in five dimensions	3
2.1 Spinor conventions in five dimensions	3
2.2 The Weyl spinor	4
2.3 The Petrov classification	4
2.4 Examples	5
3. Metrics of type <u>22</u>: equations	6
3.1 Ansatz	7
3.2 Field equations	7
3.3 The Petrov condition	8
4. Classification of metrics of type <u>22</u>	9
5. Classification of metrics of type <u>22</u>	10
6. Study of the black hole on a cylinder	10
7. Conclusions	11
A. Tetrad transformation on the Weyl spinor	12
B. The field equations	12
C. Type <u>22</u>: solutions of the field equations	14
D. Type <u>22</u>: solutions of the field equations in case A, B and C	15
E. Type <u>22</u>: solutions in case D	18

1. Introduction

In this article, we try to construct the metric of a neutral black hole in a space with one compact dimension. More specifically, we look for a black hole in a 5-dimensional spacetime which is asymptotically the product of the 4-dimensional Minkowski space and a circle. Henceforth, we will refer to this kind of black hole as *a black hole on a cylinder*.

The Newton potential of a point mass on a cylinder is easily found. Indeed, because of the linearity of the Laplace operator, one can determine this potential by a summation

over an infinite number of mirror masses. However, the Einstein equations are non-linear and, hence, determining the metric of a (neutral) black hole on a cylinder is much more difficult¹. Its metric is presumably complicated because neither the spherical symmetry nor the cylindrical symmetry applies in general. Therefore, the metric depends on two coordinates – on the radial distance in the large dimensions and on the coordinate in the compact dimension – and the Einstein equations lead to a system of partial differential equations.

Instead of a direct attempt to integrate these complicated equations, our line of attack will be the following. First, we determine the Petrov type of the 5-dimensional Schwarzschild black hole and of the black string wrapping the compact circle. These two geometries can be viewed as limiting cases of the black hole on the cylinder. Indeed, very close to the black hole on the cylinder, the size of the compact circle is irrelevant and the metric approaches the 5-dimensional Schwarzschild geometry. On the other hand, far from the black hole on the cylinder, the fact that the black hole is localized is unimportant and the metric approaches the one of the wrapped black string. It will turn out that the 5-dimensional black hole has Petrov type 22 and that the wrapped string has Petrov type 22 (the notation will be explained in section 2). Because the metrics of type 22 form a subset of the metrics of type 22, the smallest special class containing both limiting cases has Petrov type 22. Therefore, we know that if the black hole on a cylinder is algebraically special, it should have Petrov type 22.

Secondly, we classify Ricci-flat metrics that are static, have an $SO(3)$ isometry group and are in addition of Petrov type 22. It is the latter constraint which makes the Einstein equations much more tractable.

Thirdly, it turns out that the black hole on a cylinder does not show up in this classification and hence is not algebraically special².

Our method is analogous to the method used for constructing the rotating black hole in four dimensions. This metric was found by Kerr [1] while sieving³ through the metrics having 4-dimensional Petrov type D .

The rest of this article is organized as follows. In section 2, we give an outline of the 5-dimensional Petrov classification. Although there exist good textbooks [2] on the Petrov classification in 4 dimensions, the generalization to 5 dimensions of this classification scheme has to our knowledge not appeared in the literature. In section 2, we also determine the Petrov type of the 5-dimensional Schwarzschild black hole and of the black string wrapping the compact circle. In section 3, we give an overview of the equations one has to solve to find static black holes that have an $SO(3)$ isometry group and have Petrov type 22. In

¹The metric of a maximally charged black hole on a cylinder is known [3]. In this case, because there is no force between extremal black holes, the non-linearities in the Einstein equations are less severe and hence easier to solve. One can also consider this simplification as a consequence of supersymmetry.

²One could suspect that the 5-dimensional black hole and the wrapped string are algebraically special because they both have more symmetries than the black hole on a cylinder; the isometry group of the space-like slices is $SO(4)$ and $SO(3) \times U(1)$ respectively versus only $SO(3)$. However, there is no clear connection between metrics being algebraically special and the existence of Killing vectors and so this argument is not valid.

³See [4] or [5], page 265, for an exhaustive list of all 4-dimensional spacetimes of Petrov type D .

section 4, we classify the metrics that are static, have an $SO(3)$ isometry group and are of Petrov type 22. In section 5, we give a classification of metrics that are static, have an $SO(3)$ isometry group and are of Petrov type 22. In section 6, we prove that all metrics of type 22 that approach the wrapped string geometry at infinity, are homogeneous along the circle. This proves in particular that the black hole on a cylinder is not algebraically special.

We end with conclusions in section 7. Most calculations have been referred to the appendices.

2. Petrov classification in five dimensions

The Petrov type of a metric is determined by its Weyl tensor C_{ijkl} . Originally, this classification in four dimensions was done using the eigenspace structure and eigenvalues of the symmetric 6×6 matrix $C_{[ij][kl]}$. Imposing the Bianchi identity $C_{i[jkl]} = 0$ is unnatural when using this two-index notation and the classification becomes rather cumbersome. Thereafter, it was realized that the 4-dimensional Petrov classification is much easier if one uses spinors. One can define a spinor equivalent to the Weyl tensor, called the Weyl spinor Ψ_{abcd} . Instead of classifying the possible Weyl tensors, one classifies the possible Weyl spinors. In this way, the Petrov classification becomes very clear and elegant.

It turns out that in five dimensions also, it is very natural to use spinors to discuss the Petrov classification. Therefore, we will now give a brief overview of spinor conventions in five dimensions. For a more elaborate review, we refer the reader to the literature, e.g. [6]. Our spacetimes will have Euclidean signature from now on. Although our main interest lies in the construction of a *Minkowski* black hole on a cylinder, this black hole is static and can easily be Wick-rotated.

2.1 Spinor conventions in five dimensions

It is easy to find a realization of the Clifford algebra in 5 Euclidean dimensions

$$\{\gamma_i, \gamma_j\} = 2\delta_{ij}.$$

When we calculate the Petrov type of some concrete metrics later on, we will always make the following choice

$$\gamma_1 = \sigma_1 \otimes 1, \quad \gamma_2 = \sigma_2 \otimes 1, \quad \gamma_3 = \sigma_3 \otimes \sigma_1, \quad \gamma_4 = \sigma_3 \otimes \sigma_2, \quad \gamma_5 = \sigma_3 \otimes \sigma_3. \quad (2.1)$$

We use letters at the beginning of the alphabet ($a, b, c, \dots$) to denote spinor indices and letters in the middle ($i, j, k, \dots$) to denote spacetime indices. If we let the γ -matrices act on spinors ψ^a on the left, we see that they have the following index structure

$$\gamma_i \equiv (\gamma_i)^a_b.$$

We can then lower the spinor indices on the γ -matrices as follows

$$(\gamma_i)_{ab} = C_{ac}(\gamma_i)^c_b,$$

where we take the charge conjugation matrix to be $C = -i\sigma_1 \otimes \sigma_2$. The matrix $(\gamma_i)_{ab}$ is antisymmetric. We also have

$$(\gamma_{ij})_{ab} = C_{ac}(\gamma_{ij})^c_b \quad \text{where } \gamma_{ij} = \frac{1}{2}[\gamma_i, \gamma_j].$$

This matrix is symmetric. We will use it to define the Weyl spinor in the next section.

2.2 The Weyl spinor

In four dimensions, the definition of the Weyl spinor is motivated by the isometry between $so(4)$ and $su(2) \oplus su(2)$. Similarly, in five dimensions, we use the isomorphism between $so(5)$ and $sp(4)$ to map representations of $so(5)$ into representations of $sp(4)$ and vice versa. The Weyl tensor C_{ijkl} of $so(5)$ is mapped into a tensor Ψ_{abcd} of $sp(4)$, which, using the Bianchi identity, turns out to be completely symmetric

$$so(5) \rightarrow sp(4) : \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \rightarrow \square\square\square\square : \Psi_{abcd} = (\gamma_{ij})_{ab}(\gamma_{kl})_{cd}C^{ijkl}. \quad (2.2)$$

We will refer to the object Ψ_{abcd} as the *Weyl spinor*. As an easy check for this correspondence, one can count the number of degrees of freedom of the completely symmetric tensor Ψ_{abcd} . This is 35, equal to the dimension of the Weyl tensor C_{ijkl} . It is easy to invert the relation (2.2):

$$C_{ijkl} = \frac{1}{64}(\gamma_{ij})^{ab}(\gamma_{kl})^{cd}\Psi_{abcd}. \quad (2.3)$$

2.3 The Petrov classification in five dimensions

The fact that we have found a completely symmetric spinor equivalent of the Weyl tensor makes it easy to classify Weyl tensors in an invariant way. To this end, we associate to the Weyl spinor Ψ_{abcd} the *Weyl polynomial* Ψ , which is a homogeneous polynomial of degree four in four variables:

$$\Psi = \Psi_{abcd}x^ax^bx^cx^d. \quad (2.4)$$

The Petrov type of a given Weyl tensor is nothing else than the number and multiplicity of irreducible⁴ factors of Ψ . This is an invariant classification scheme, because under $O(5)$ transformations of the tetrad, the Weyl polynomial changes by a linear transformation of its variables x^a . This transformation respects the factorization properties of Ψ . In this way, we get 12 different Petrov types, which are depicted in figure 1. We use the following notation. The number denotes the degree of the irreducible factors and underbars denote the multiplicities. For example, a Weyl polynomial which can be factorized into two different factors, each having degree 2, has Petrov type $\underline{2}\underline{2}$. As a second example, a Weyl polynomial which can be factorized in a polynomial of degree 2 times the square of

⁴The Weyl polynomial has generically complex coefficients. Hence, in this article, we will decompose this Weyl polynomial over $\mathbb{C}$. However, the Weyl polynomial is in fact a real polynomial in the sense that there is a reality condition on the Weyl spinor Ψ_{abcd} derived from the fact that the γ -matrices are Hermitian. Therefore, a more refined classification is possible if one makes a distinction between “real” and “complex” factorizations. Furthermore, it is possible to refine the Petrov classification by characterizing the leftover polynomials of degree 2 by their eigenvalues, for example.




Figure 1: The 12 different Petrov types in 5 dimensions.

a polynomial of degree 1 has Petrov type $\underline{211}$. The arrows between the different Petrov types denote increasing specialization of the Weyl tensor. In accord with the literature on the 4-dimensional Petrov classification, all Weyl tensors different from type 4 are called algebraically special.

2.4 Examples

five-dimensional spherical symmetric black hole

The metric reads

$$ds^2 = (1 + 2V)dt^2 + (1 + 2V)^{-1}dr^2 + r^2 d\Omega_3^2, \quad (2.5)$$

where the function $V(r)$ is the Newton potential $V(r) = -G_5 M/r^2$ and $d\Omega_3^2$ is the metric on the unit 3-sphere

$$d\Omega_3^2 = d\psi^2 + \sin^2 \psi (d\theta^2 + \sin^2 \theta d\phi^2).$$

If we choose the tetrad as

$$\begin{aligned} e_t &= \frac{1}{\sqrt{1 + 2V}} \partial_t & e_r &= \sqrt{1 + 2V} \partial_r & e_\psi &= \frac{1}{r} \partial_\psi \\ e_\theta &= \frac{1}{r \sin \psi} \partial_\theta & e_\phi &= \frac{1}{r \sin \psi \sin \theta} \partial_\phi, \end{aligned}$$

we find after some algebra the Weyl polynomial

$$\Psi = -\frac{192 G_5 M}{r^4} (xt - yz)^2.$$

Here we have used the notation $(x^1, x^2, x^3, x^4) = (x, y, z, t)$. Hence, the metric (2.5) has Petrov type $\underline{22}$.

Wrapped black string

The metric of a black string wrapped around a circle of radius R reads

$$ds^2 = (1 + 2V)dt^2 + (1 + 2V)^{-1}dr^2 + r^2 d\Omega_2^2 + R^2 d\alpha^2, \quad (2.6)$$

where $V(r)$ is the Newton potential $V(r) = -G_4 M/r$ and $d\Omega_2^2$ is the metric on the unit 2-sphere. If we choose the tetrad as

$$\begin{aligned} e_t &= \frac{1}{\sqrt{1+2V}} \partial_t & e_r &= \sqrt{1+2V} \partial_r & e_\alpha &= \frac{1}{R} \partial_\alpha \\ e_\theta &= \frac{1}{r} \partial_\theta & e_\phi &= \frac{1}{r \sin \theta} \partial_\phi, \end{aligned}$$

we find after some algebra the Weyl polynomial

$$\Psi = -\frac{96 G_4 M}{r^3} (x^2 t^2 + y^2 z^2).$$

The metric (2.6) has Petrov type 22 because we can factorize Ψ as

$$\Psi = -\frac{96 G_4 M}{r^3} (xt + iyz) (xt - iyz).$$

At this point, we know the Petrov type of two limiting cases of the black hole on a cylinder. Namely, at large distances (or, equivalently, at small compactification scale), the metric has Petrov type 22. At small distances (or, equivalently, large compactification scale), the metric has Petrov type 22. Taking a look at figure 1, we see that the black hole on a cylinder should have Petrov type 22 or type 4. In the latter case, the metric is algebraically generic. In the next section, we give the equations one has to solve to determine all metrics that are static, have an $SO(3)$ isometry group and are of Petrov type 22.

3. Metrics of type 22: equations

Because we want to impose a constraint on the Weyl tensor, we will not solve the Einstein equations in the usual way. We will not write an ansatz for the metric and solve the differential equations resulting from putting the Ricci tensor to zero. Instead, we will solve the following set of equations for the dual tetrad θ , the connection ω and the curvature Ω

$$d\theta + \omega \wedge \theta = 0 \tag{3.1a}$$

$$d\omega + \omega \wedge \omega = \Omega \tag{3.1b}$$

$$d\Omega + \omega \wedge \Omega - \Omega \wedge \omega = 0. \tag{3.1c}$$

These three equations are of course respectively the definition of zero torsion, the definition of the curvature Ω and the Bianchi identity. In the last two equations (3.1b, 3.1c), we use equation (2.3) to express the Riemann tensor Ω in terms of the Weyl spinor Ψ_{abcd} . We can indeed use formula (2.3) for this substitution because the Riemann tensor is equal to the Weyl spinor for spacetimes with zero Ricci tensor. Because the Weyl spinor now enters the field equations explicitly, it is easy to impose a condition on the Petrov type.

3.1 Ansatz

The most general ansatz for the tetrad of a static spacetime with $SO(3)$ symmetry reads

$$\begin{aligned} e_t &= A \partial_t \\ e_r \text{ and } e_\alpha &: \text{linear combinations of } \partial_r \text{ and } \partial_\alpha \\ e_\theta &= \sigma \partial_\theta \\ e_\phi &= \frac{\sigma}{\sin \theta} \partial_\phi \end{aligned} \tag{3.2}$$

Here, A and σ are functions of the coordinates r and α . The most general connection consistent with the above ansatz is

$$\omega = \begin{pmatrix} 0 & \mu\theta^1 & \nu\theta^1 & 0 & 0 \\ -\mu\theta^1 & 0 & \delta\theta^2 + \varepsilon\theta^3 & \kappa\theta^4 & \kappa\theta^5 \\ -\nu\theta^1 & -\delta\theta^2 - \varepsilon\theta^3 & 0 & \lambda\theta^4 & \lambda\theta^5 \\ 0 & -\kappa\theta^4 & -\lambda\theta^4 & 0 & -\sigma \cot \theta \theta^5 \\ 0 & -\kappa\theta^5 & -\lambda\theta^5 & \sigma \cot \theta \theta^5 & 0 \end{pmatrix}. \tag{3.3}$$

Here, $(\theta^1, \dots, \theta^5)$ is the basis dual to the tetrad (3.2) and the seven functions $(\mu, \nu, \delta, \varepsilon, \kappa, \lambda, \sigma)$ are functions of the coordinates r and α . Notice that although we could have expressed the connection ω in terms of the functions appearing in the tetrad (3.2), we have not done so. Rather, as will become clear in the following, we determine ω by solving the equations (3.1b) and (3.1c). Finally, a calculation shows that the most general ansatz for the Weyl spinor Ψ_{abcd} has 4 independent components.

$$\begin{aligned} \Psi_{1114} &= \Psi_{1444} = \Psi_{2223} = \Psi_{2333} = -3\psi_1 \\ \Psi_{1123} &= \Psi_{1224} = \Psi_{1334} = \Psi_{2344} = \psi_1 \\ \Psi_{1111} &= \Psi_{2222} = \Psi_{3333} = \Psi_{4444} = -3\psi_2 \\ \Psi_{1133} &= \Psi_{2244} = \psi_2 \\ \Psi_{2233} &= \Psi_{1144} = \psi_3 \\ \Psi_{1122} &= \Psi_{3344} = \psi_4 \text{ and } \Psi_{1234} = -\frac{1}{2}(\psi_3 + \psi_4) \end{aligned} \tag{3.4}$$

Here again, the four functions ψ_i are functions of the coordinates r and α . In some cases, equations simplify when written in terms of the functions φ_i , defined by

$$\begin{aligned} \psi_1 &= -8i\varphi_4 & \psi_2 &= 8(\varphi_2 + \varphi_3) \\ \psi_3 &= 8(\varphi_2 - 3\varphi_3) & \psi_4 &= 8(-2\varphi_1 - \varphi_2 + \varphi_3). \end{aligned} \tag{3.5}$$

The above linear combinations are chosen in such a way that the functions φ_i transform nicely under tetrad transformations (see appendix A).

3.2 Field equations

We can now take the ansätze (3.3) and (3.4) and insert them into equations (3.1). The differential equations obtained in this way are quite lengthy. They can be found in appendix B.

3.3 The Petrov condition

In this section, we give necessary and sufficient conditions on the components φ_i of the Weyl spinor such that the Weyl polynomial Ψ obtained from the Weyl spinor (3.4) has Petrov type 22.

The metric has Petrov type 22 if and only if one of the following four conditions is satisfied

$$\begin{array}{ll} \text{Case A: } \varphi_3 = \varphi_4 = 0 & \text{Case C: } \varphi_1^2 = \varphi_3^2 + \varphi_4^2 \\ \text{Case B: } \varphi_1 + \varphi_2 = 0 & \text{Case D: } \varphi_2^2 = \varphi_3^2 + \varphi_4^2 \end{array} \quad (3.6)$$

The characterization is most easily derived in a frame for which $\varphi_4 = 0$. It is proven in appendix A that such a frame exists. In this frame, using (3.4) and (3.5) and dropping an irrelevant overall constant, the Weyl polynomial reads

$$\begin{aligned} \Psi_{\text{fr}} = & \psi_2 ((x^2 - z^2)^2 + (y^2 - t^2)^2) - 2\psi_3(x^2 t^2 + y^2 z^2) \\ & - 2\psi_4(x^2 y^2 + z^2 t^2) + 4(\psi_3 + \psi_4) x y z t . \end{aligned} \quad (3.7)$$

The subscript of Ψ_{fr} is to remind the reader that we have chosen a particular frame. We want the polynomial (3.7) to be a product of 2 factors of degree 2. We can deduce necessary conditions for this reducibility in the following way. The polynomial (3.7) should still be reducible when we put one of the variables x, y, z, t to zero. When we restrict Ψ_{fr} to only three variables in this manner, the algebra becomes less cumbersome and one can show that the restricted Ψ_{fr} can be factorized into 2 factors of degree 2 if and only if one of the following four cases holds

$$\psi_2 = 0 \quad \text{or} \quad \psi_2^2 = \psi_3^2 \quad \text{or} \quad \psi_2^2 = \psi_4^2 \quad \text{or} \quad \psi_3 + \psi_4 = 0. \quad (3.8)$$

Although these conditions were derived as necessary conditions for having Petrov type 22, it turns out that they are sufficient as well. In the first three cases, the polynomial Ψ_{fr} is easily factorized after writing it as a sum of squares

$$\begin{array}{ll} \text{if } \psi_2 = 0, & \text{then } \Psi_{\text{fr}} = -2(\psi_3(xt - yz)^2 + \psi_4(xy - zt)^2) \\ \text{if } \psi_2^2 = \psi_3^2, & \text{then } \Psi_{\text{fr}} = \psi_2(x^2 \pm y^2 - z^2 \mp t^2)^2 - 2(\psi_3 + \psi_4)(xy - zt)^2 \\ \text{if } \psi_2^2 = \psi_4^2, & \text{then } \Psi_{\text{fr}} = \psi_2(x^2 \pm y^2 - z^2 \mp t^2)^2 - 2(\psi_3 + \psi_4)(xt - yz)^2 \end{array}$$

The factorization in the fourth case is straightforward. We give it for completeness.

$$\begin{aligned} \text{if } \psi_3 + \psi_4 = 0, \quad \text{then } \Psi_{\text{fr}} = & \frac{1}{\psi_2} (\psi_2 x^2 + (\psi_3 + w)y^2 - \psi_2 z^2 + (\psi_4 - w)t^2) \times \\ & \times (\psi_2 x^2 + (\psi_3 - w)y^2 - \psi_2 z^2 + (\psi_4 + w)t^2), \end{aligned}$$

where $w^2 = \psi_4^2 - \psi_2^2$. The conditions (3.8) are expressed in a frame in which $\psi_1 = 0$. Rotating this frame back to its general position leads directly to the conditions (3.6).

4. Classification of metrics of type 22

Besides the five-dimensional black hole and the wrapped black string, both treated in section 2.4, we have encountered the following metrics:

1. the Kasner metric with exponents p_i

$$ds^2 = dt^2 + \sum_{i=1}^4 \left(\frac{t}{t_0} \right)^{2p_i} (dx^i)^2. \quad (4.1)$$

This metric is Ricci flat if $\sum_i p_i = \sum_i p_i^2 = 1$. It appears naturally in the classification because, as is well known, Kasner spaces describe the near-singularity geometry of Schwarzschild spaces.

2. the cone on a dS-Schwarzschild space

$$ds^2 = dt^2 + t^2 \left[\left(1 - r^2 + \frac{M}{r} \right) d\alpha^2 + \frac{dr^2}{1 - r^2 + \frac{M}{r}} + r^2 d\Omega^2 \right] \quad (4.2)$$

3. a vacuum cosmological model in which space is the product of two spheres

$$ds^2 = dt^2 + a(t)^2 d\Omega^2 + b(t)^2 d\Omega^2 \quad (4.3)$$

Here, the functions $a(t)$ and $b(t)$ are determined by the vacuum Einstein equations. We do not know the analytical expression of the general solution. A particular solution is the cone on the product of two spheres $ds^2 = dt^2 + t^2/3 (d\Omega^2 + d\Omega^2)$

4. a vacuum cosmological model on the product of a sphere and the hyperbolic plane $\mathbb{H}$

$$ds^2 = dt^2 + a(t)^2 d\Omega^2 + b(t)^2 d\mathbb{H}^2 \quad (4.4)$$

5. a vacuum cosmological model in which space is the product of a sphere and a plane

$$ds^2 = -4 \cosh^{-6}(t/t_0) \sinh^{2-\frac{4}{\sqrt{3}}}(t/t_0) dt^2 + \sinh^{\frac{2}{\sqrt{3}}}(t/t_0) (dx^2 + dy^2) + t_0^2 \cosh^{-4}(t/t_0) \sinh^{2-\frac{4}{\sqrt{3}}}(t/t_0) d\Omega^2 \quad (4.5)$$

6. the metric of another homogeneous wrapped object

$$ds^2 = (1 - 2M/r) dt^2 + dr^2 + R^2 (1 - 2M/r)^{-1} d\alpha^2 + r^2 (1 - 2M/r) d\Omega^2 \quad (4.6)$$

This metric has curvature singularities in $r = 0$ and $r = 2M$.

Some of these solutions admit 3 orthogonal commuting Killing vectors and therefore belong to the family of higher dimensional Weyl solutions that Reall and Emparan studied [17]. In particular, the five-dimensional black hole, the black string and the solutions (4.1), (4.5), and (4.6) all belong to this family.

The metrics (2.6), (4.5) and (4.6) are all special cases of the general homogeneous solution [18]

$$ds^2 = \left(\frac{r-m}{r+m} \right)^{2a} dt^2 + \frac{1}{r^4} (r+m)^{2+2a+2c} (r-m)^{2-2a-2c} (dr^2 + r^2 d\Omega^2) + \left(\frac{r-m}{r+m} \right)^{2c} d\alpha^2$$

with $a^2 + ac + c^2 = 1$. In particular, for the wrapped string (2.6), $a = 1$ and $c = 0$, for the metric (4.6), $a = 1$ and $c = -1$ and for the metric (4.5), $a = c = 1/\sqrt{3}$. The classification is then as follows⁵.

Case A: the wrapped string (2.6)

Case B: the five-dimensional black hole (2.5)

Case C: the five-dimensional black hole (2.5) and the metric (4.6)

Case D: the 5-dimensional black hole, the cone on an dS-Schwarzschild space (4.2), the wrapped string and the cosmological models (4.3), (4.4) and (4.5)

The derivation of this classification is tedious, it can be found in the appendices D and E.

5. Classification of metrics of type 22

From the characterization (3.6), one can easily derive the necessary and sufficient conditions leading to metrics of Petrov type 22. These conditions are combinations of the conditions (3.6), which is reflected in our notation. Details of the classification can be found in appendix C.

The metric is of Petrov type 22 in the following three cases:

Case AC: $\varphi_1 = \varphi_3 = \varphi_4 = 0$: only flat space

Case AD: $\varphi_2 = \varphi_3 = \varphi_4 = 0$: only flat space

Case BC: $\varphi_1 + \varphi_2 = 0$ and $\varphi_1^2 = \varphi_3^2 + \varphi_4^2$: the five-dimensional black hole (2.5)

6. Study of the black hole on a cylinder

We now prove that the black hole on a cylinder and – more generally – all inhomogeneous wrapped strings are not algebraically special.

These metrics approach the geometry of the wrapped string at infinity, hence, we see from figure 1 that they should be of Petrov type 22 if they are algebraically special.

⁵For brevity, we have adopted the following usage. For example, we say that case B consists of the five-dimensional Schwarzschild metric. It is to be understood that its two limits – flat space and the Kasner space (4.1) with exponents $(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ – belong to case B as well.

However, in section 4, we have classified all solutions that are static, have an $SO(3)$ isometry group and have Petrov type 22. The non-homogeneous wrapped string was not one of them. Hence, we know that the black hole on a cylinder and more generally the non-homogeneous wrapped string are not algebraically special. This completes the proof.

7. Conclusions

It is well-known that string theory suggests that spacetime has 10 or 11 dimensions. Although string theory is not sufficiently developed to make concrete physical predictions that can be tested experimentally, it seems to be clear that these extra dimensions should be very small for string theory not to be in disagreement with reality⁶.

It is interesting to study black holes in these higher dimensional spacetimes for the following reasons. Firstly, due to the Einstein equations, black holes will alter the geometry of these extra dimensions in their neighborhood. It is possible that this will lead to an increase in size of these extra dimensions close to a black hole, making these dimensions easier to detect. Secondly, string theory predicts generically that the fundamental physical constants like the gravitational constant, masses of elementary particles and the fine structure constant should depend on the geometry of the compactification space. Therefore, knowledge of the black hole geometry can lead to some predictions of the behaviour of these constants near black holes.

Thirdly, Gregory and Laflamme [8] found that a neutral black string wrapping a cylinder is classically unstable if its mass is sufficiently small. Therefore, it seems natural to conjecture that these black strings will decay into a black hole that is localized in the compact direction. However, recently, Horowitz and Maeda [9] argued that, classically, the black string would not decay into the black hole on a cylinder but rather to a *non-homogeneous* wrapped string⁷. A similar claim holds for black strings with large masses: in [12], it was argued that black strings with large masses have smaller entropy than the non-homogeneous wrapped strings, and hence should decay as well.

In this article, we studied the metrics of the black hole on a cylinder and of the non-homogeneous wrapped string using the notion of algebraically special metrics. We have treated the Petrov classification in five dimensions. We have given a complete classification of five-dimensional metrics that are Ricci-flat, static, of Petrov type 22 and have an $SO(3)$ isometry group. We used this classification to prove that the black hole on a cylinder and the non-homogeneous wrapped string are not algebraically special.

Some possible topics for future research are

1. Classify all Ricci-flat metrics of type 22 without the additional assumptions we made. This could lead to the discovery of new five-dimensional metrics. One knows that the set of black holes is much richer in five dimensions than in four dimensions. See [13] for a study of uniqueness theorems in higher dimensions and [14] for a five-dimensional rotating black string.

⁶Recently, however, scenarios with large extra dimensions have been proposed [7].

⁷See [10] for a nice review and [11] for numerical investigations on these non-homogeneous black strings.

2. The metric of a black hole in the RS-scenario [7] that has a localized horizon in the bulk spacetime is not known [15, 16]. It is possible that a line of research similar to the one presented here would lead to a construction of this metric.
3. It would be nice to find some other simplifying assumption yielding solvable equations. It may be possible to find the metric of the black hole on a cylinder by a clever choice of coordinates; for an attempt, see [12].

Acknowledgments

I would like to thank Yves Demasure, Stefan Vandoren, Antoine Van Proeyen and Walter Troost for discussions. I also would like to thank Pierre Savaria and Walter Troost for suggestions to improve this text and S. Hervik for a correction in formula (4.5) in an earlier version of this paper. This work was supported in part by the European Commission RTN program HPRN-CT-2000-00131.

A. Tetrad transformation on the Weyl spinor

In this appendix, we give the behaviour of the components of the Weyl spinor under tetrad transformations. Direct use of the definition (2.2), shows that under tetrad rotations ($s = +1$) and reflections ($s = -1$)

$$\begin{pmatrix} e_r \\ e_\alpha \end{pmatrix} \rightarrow \begin{pmatrix} \cos \chi & \sin \chi \\ -s \sin \chi & s \cos \chi \end{pmatrix} \begin{pmatrix} e_r \\ e_\alpha \end{pmatrix},$$

the Weyl spinor transforms as

$$\varphi_1 \rightarrow \varphi_1, \quad \varphi_2 \rightarrow \varphi_2, \quad \begin{pmatrix} \varphi_3 \\ \varphi_4 \end{pmatrix} \rightarrow \begin{pmatrix} \cos 2\chi & -\sin 2\chi \\ s \sin 2\chi & s \cos 2\chi \end{pmatrix} \begin{pmatrix} \varphi_3 \\ \varphi_4 \end{pmatrix}.$$

From this, we have the important result that we can always use a tetrad rotation to put $\varphi_4 = 0$. On top of this, we can use a reflection to specify the sign of φ_3 .

B. The field equations

In this appendix, we explicitly write down the equations resulting from inserting the static axi-symmetric ansatz (3.2, 3.3, 3.4) into the differential equations (3.1). We will split up these equations into three blocks.

The equations from (3.1a)

The condition of zero torsion (3.1a) yields the commutation relations between the tetrad vectors (3.2). The non-zero commutators are

$$[e_t, e_r] = \mu e_t, \quad [e_t, e_\alpha] = \nu e_t, \quad (\text{B.1a})$$

$$[e_r, e_\theta] = \kappa e_\theta, \quad [e_\alpha, e_\theta] = \lambda e_\theta, \quad (\text{B.1b})$$

$$[e_r, e_\phi] = \kappa e_\phi, \quad [e_\alpha, e_\phi] = \lambda e_\phi, \quad (\text{B.1c})$$

$$[e_r, e_\alpha] = \delta e_r + \varepsilon e_\alpha \quad (\text{B.1d})$$

$$[e_\theta, e_\phi] = -\cot \theta \sigma e_\phi. \quad (\text{B.1e})$$

The equations from (3.1b)

In the definition (3.1b) of the curvature tensor Ω , we use equation (2.3) and the fact that the Ricci tensor is zero to express the curvature tensor in terms of the Weyl spinor Ψ_{abcd} . We also use equation (3.1a) to replace terms involving $d\theta^i$ that appear in $d\omega$. In this way, we find two algebraic equations

$$\sigma^2 - \kappa^2 - \lambda^2 - 3\varphi_1 - \varphi_2 = 0 \quad (\text{B.2})$$

$$\kappa\mu + \lambda\nu + \varphi_1 - \varphi_2 = 0 \quad (\text{B.3})$$

and a set of first order differential equations. The ones involving the vector e_r are

$$e_r(\sigma) = \sigma\kappa \quad (\text{B.4a})$$

$$e_r(\varepsilon) - e_\alpha(\delta) = \delta^2 + \varepsilon^2 + \varphi_1 + 3\varphi_2 \quad (\text{B.4b})$$

$$e_r(\kappa) = \kappa^2 - \delta\lambda - \varphi_1 - \varphi_2 - 2\varphi_3 \quad (\text{B.4c})$$

$$e_r(\lambda) = \kappa\delta + \kappa\lambda + 2\varphi_4 \quad (\text{B.4d})$$

$$e_r(\mu) = -\nu\delta - \mu^2 - \varphi_1 + \varphi_2 - 4\varphi_3 \quad (\text{B.4e})$$

$$e_r(\nu) = -\nu\mu + \delta\mu + 4\varphi_4 \quad (\text{B.4f})$$

and the ones involving the vector e_α are

$$e_\alpha(\sigma) = \sigma\lambda \quad (\text{B.5a})$$

$$e_\alpha(\kappa) = -\varepsilon\lambda + \kappa\lambda + 2\varphi_4 \quad (\text{B.5b})$$

$$e_\alpha(\lambda) = \varepsilon\kappa + \lambda^2 - \varphi_1 - \varphi_2 + 2\varphi_3 \quad (\text{B.5c})$$

$$e_\alpha(\mu) = -\nu\varepsilon - \nu\mu + 4\varphi_4 \quad (\text{B.5d})$$

$$e_\alpha(\nu) = -\nu^2 + \varepsilon\mu - \varphi_1 + \varphi_2 + 4\varphi_3 \quad (\text{B.5e})$$

The equations from (3.1c)

These are the equations resulting from the Bianchi identity. We again replace the curvature tensor Ω by the Weyl spinor Ψ_{abcd} . We also use equation (3.1a) to replace terms involving

$d\theta^i$ that appear in $d\Omega$. In this way, we find the following set of equations

$$e_r(\varphi_1) = \frac{5}{2}\kappa \varphi_1 + \frac{1}{2}(\kappa + \mu)\varphi_2 + \frac{1}{2}(4\kappa + \mu)\varphi_3 - \frac{1}{2}(\nu + 4\lambda)\varphi_4 \quad (\text{B.6a})$$

$$e_r(\varphi_2) = \frac{1}{2}\kappa \varphi_1 + \frac{1}{2}(5\kappa - 3\mu)\varphi_2 - \frac{1}{2}(4\kappa + 3\mu)\varphi_3 + \frac{1}{2}(3\nu + 4\lambda)\varphi_4 \quad (\text{B.6b})$$

$$e_r(\varphi_3) = e_\alpha(\varphi_4) + \frac{1}{2}\kappa\varphi_1 - \frac{1}{2}(\kappa + \mu)\varphi_2 + (2\varepsilon + \kappa - \frac{1}{2}\mu)\varphi_3 \quad (\text{B.6c})$$

$$+ (\frac{1}{2}\nu + 2\delta - \lambda)\varphi_4 \quad (\text{B.6d})$$

$$e_r(\varphi_4) = -e_\alpha(\varphi_3) - \frac{1}{2}\lambda \varphi_1 + \frac{1}{2}(\nu + \lambda)\varphi_2 + (-\frac{1}{2}\nu - 2\delta + \lambda)\varphi_3 \quad (\text{B.6e})$$

$$+ (2\varepsilon + \kappa - \frac{1}{2}\mu)\varphi_4 \quad (\text{B.6f})$$

$$e_\alpha(\varphi_1) = \frac{5}{2}\lambda \varphi_1 + \frac{1}{2}(\nu + \lambda)\varphi_2 - \frac{1}{2}(\nu + 4\lambda)\varphi_3 - \frac{1}{2}(4\kappa + \mu)\varphi_4 \quad (\text{B.6g})$$

$$e_\alpha(\varphi_2) = \frac{1}{2}\lambda\varphi_1 + \frac{1}{2}(-3\nu + 5\lambda)\varphi_2 + \frac{1}{2}(3\nu + 4\lambda)\varphi_3 + \frac{1}{2}(4\kappa + 3\mu)\varphi_4 \quad (\text{B.6h})$$

C. Type 22: solutions of the field equations

In this appendix, we classify all solutions of the field equations given in appendix B that are of Petrov type 22.

Case AC: $\varphi_1 = \varphi_3 = \varphi_4 = 0$

From the Bianchi equations (B.6), we immediately get $(\kappa + \mu)\varphi_2 = 0$ and $(\lambda + \nu)\varphi_2 = 0$. Hence, we have two possibilities. The first one is $\varphi_2 = 0$. Then all components of the Weyl spinor are zero and we obtain flat space. The second possibility is $\kappa + \mu = 0$ and $\lambda + \nu = 0$. Then we obtain from (B.2) and (B.3) $\sigma = 0$. This is not a good solution of the field equations, because in this case the tetrad (3.2) is degenerate. Therefore, in case AC, we obtain only flat space.

Case AD: $\varphi_2 = \varphi_3 = \varphi_4 = 0$

This case is similar to the previous one. From the Bianchi equations (B.6), we get $\kappa\varphi_1 = 0$ and $\lambda\varphi_1 = 0$. If $\varphi_1 = 0$, we have flat space. If $\kappa = \lambda = 0$, we find from equation (B.3) $\varphi_1 = 0$, giving flat space again.

Case BC: $\varphi_1 + \varphi_2 = 0$ and $\varphi_1^2 = \varphi_3^2 + \varphi_4^2$

As shown in appendix A, we can choose our tetrad in such a way that $\varphi_4 = 0$ and $\varphi_3 = \varphi_1$. From the Bianchi equations (B.6), we obtain 3 algebraic equations

$$\nu\varphi_1 = \delta\varphi_1 = (\kappa - \varepsilon)\varphi_1 = 0.$$

From these equations, we get either $\varphi_1 = 0$ or $\nu = \delta = \kappa - \varepsilon = 0$. The first possibility leads to flat space, hence, we continue with the second possibility. From equation (B.3), we get $\varphi_1 = -1/2 \kappa\mu$. Inserting all this into equation (B.4) and (B.5) leads to the following equations

$$e_r(\kappa) = \kappa^2 + \kappa\mu \quad (\text{C.1a}) \quad e_\alpha(\kappa) = 0 \quad (\text{C.1d})$$

$$e_r(\lambda) = \kappa\lambda \quad (\text{C.1b}) \quad e_\alpha(\lambda) = \kappa^2 + \lambda^2 - \kappa\mu \quad (\text{C.1e})$$

$$e_r(\mu) = 3\kappa\mu - \mu^2 \quad (\text{C.1c}) \quad e_\alpha(\mu) = 0 \quad (\text{C.1f})$$

We will assume from now on that $\kappa \neq 0$ and $\mu \neq 0$, because otherwise, we find flat space again. We choose our coordinates in such a way that $e_r = A\partial_r$ and $e_\alpha = B\partial_\alpha$. It then follows from (C.1d) and (C.1f) that κ and μ do not depend on the coordinate α . From the commutation relation (B.1d), we get two differential equations

$$A\partial_r B = \kappa B \quad \text{and} \quad \partial_\alpha A = 0.$$

Hence, A does not depend on r and we can put $A = -r\kappa$ by a coordinate transformation on r . The general solution for B is $B = f(\alpha)/r$, where f is an arbitrary function of α . We can subsequently do a coordinate transformation on α to put $f(\alpha) = 1$. Hence, at this point, we have $e_r = -r\kappa\partial_r$ and $e_\alpha = 1/r\partial_\alpha$. The solution of the differential equations (C.1) is now easily found

$$\kappa = -\left(\frac{C_1}{r^4} + \frac{C_2}{r^2}\right)^{1/2}, \quad \lambda = -\frac{\sqrt{C_2}}{r} \cot(\sqrt{C_2}\alpha), \quad \mu = -\frac{C_1}{r^4} \left(\frac{C_1}{r^4} + \frac{C_2}{r^2}\right)^{-1/2}.$$

Here C_1 and C_2 are two arbitrary integration constants. We have used a shift on the coordinate α to eliminate the third integration constant. Finally, the remaining tetrad vectors e_t and e_θ are determined from the commutation relation (B.1a) and the algebraic equation (B.2) respectively

$$e_t = \left(\frac{C_1}{r^2} + C_2\right)^{-1/2} \partial_t \quad \text{and} \quad \sigma = \frac{\sqrt{C_2}}{r \sin(\sqrt{C_2}\alpha)}.$$

If $C_2 \neq 0$, this solution is the 5-dimensional black hole (2.5). If $C_2 = 0$, this solution is the Kasner space (4.1) with exponents $(-\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$.

D. Type 22: solutions of the field equations in case A, B and C

In this appendix, we classify all solutions of the field equations given in appendix B, subject to the conditions of case A, B and C of (3.6).

Case A: $\varphi_3 = \varphi_4 = 0$

The Bianchi identities give

$$\kappa\varphi_1 - (\kappa + \mu)\varphi_2 = 0 \quad \text{and} \quad -\lambda\varphi_1 + (\lambda + \nu)\varphi_2 = 0. \quad (\text{D.1})$$

Because we are not interested in the trivial case $\varphi_1 = \varphi_2 = 0$, it follows from these equations that $\kappa\nu = \lambda\mu$. We choose our tetrad such that $\nu = 0$. At this point, it is necessary to resolve two cases: $\mu = 0$ and $\mu \neq 0$.

- Case A.1: $\mu = 0$

Because (μ, ν) transforms as a vector under tetrad rotations and μ and ν both are zero, we still have not fixed the tetrad. We will fix it by choosing $\lambda = 0$. From equation (B.3) we obtain $\varphi_1 = \varphi_2$. From equation (B.5c) then follows that $2\varphi_1 = \varepsilon\kappa$. Because we are not interested in the flat case $\varphi_1 = 0$, it follows that $\varepsilon \neq 0$ and $\kappa \neq 0$.

We then find from equation (B.4d) that $\delta = 0$. We will now choose our coordinates in such a way that $e_r = A\partial_r$ and $e_\alpha = B\partial_\alpha$. From equation (B.5b) and (B.6g), it follows that κ and ε do not depend on the coordinate α . From the commutation relation (B.1d), we find two differential equations

$$\partial_\alpha A = 0 \quad \text{and} \quad A\partial_r B = \varepsilon B.$$

The first of these equations tells us that A depends only on the coordinate r . We can then put $A = -r\kappa$ by a coordinate transformation on r . After these preliminary manipulations, we can start solving the differential equations. The equations (B.4) yield $e_r(\varepsilon) = \varepsilon^2 + 2\varepsilon\kappa$ and $e_r(\kappa) = \kappa^2 - \varepsilon\kappa$. The solution of these equations reads

$$\kappa = -\frac{1}{r} \left(C_1 - 2\frac{C_2}{r} \right)^{1/2} \quad \text{and} \quad \varepsilon = -\frac{C_2}{r^2} \left(C_1 - 2\frac{C_2}{r} \right)^{-1/2},$$

where C_1 and C_2 are two integration constants. The remaining tetrad vectors can now be determined from (B.1) and (B.2). They are

$$e_t = \partial_t, \quad e_\alpha = \left(C_1 - 2\frac{C_2}{r} \right)^{-1/2} \partial_\alpha, \quad \sigma = \frac{\sqrt{C_1}}{r}.$$

After changing t with α , one sees that this solution is the wrapped black string (2.6).

• Case A.2: $\mu \neq 0$

In this case, we have $\lambda = 0$. From equation (B.4f) we find $\delta = 0$, and from equations (B.3), (B.5e) and (B.5c) we obtain

$$\varepsilon = -\kappa, \quad \varphi_1 = -\frac{\kappa}{2}(\kappa + \mu) \quad \text{and} \quad \varphi_2 = \frac{\kappa}{2}(-\kappa + \mu).$$

Inserting these values into (D.1), we find $\kappa(\kappa + \mu)\mu = 0$. Hence, $\varphi_1 = 0$, and the treatment in case AC of appendix C shows that this leads to flat space.

Case B: $\varphi_1 + \varphi_2 = 0$

We choose the tetrad in such a way that $\varphi_4 = 0$. From the Bianchi equations (B.6), we get $\mu(\varphi_1 - \varphi_3) = 0$ and $\nu(\varphi_1 + \varphi_3) = 0$. Hence, there are two possibilities. The first one is $\mu \neq 0$ or $\nu \neq 0$. Then we have $\varphi_1 = \varphi_3$ or $\varphi_1 = -\varphi_3$. Therefore the metric has type 22. This case is treated in case BC of appendix C. The second possibility is $\mu = \nu = 0$. Then, it follows from (B.5e) and (B.4e) that $\varphi_1 = \varphi_2 = \varphi_3 = 0$. Hence, this possibility yields flat space.

Case C: $\varphi_1^2 = \varphi_3^2 + \varphi_4^2$

We choose our tetrad such that $\varphi_4 = 0$ and $\varphi_3 = \varphi_1$. From the Bianchi identities (B.6), we obtain two algebraic equations

$$(3\kappa - 2\varepsilon + \mu)\varphi_1 + (\kappa + \mu)\varphi_2 = 0 \quad (\text{D.2}) \quad \text{and} \quad \delta\varphi_1 = 0. \quad (\text{D.3})$$

We assume from now on that $\varphi_1 \neq 0$, because otherwise, the discussion in case AC of appendix C shows that this leads to flat space. Hence, we have $\delta = 0$. We could combine (D.2) and (B.3) to obtain rather complicated expressions for φ_1 and φ_2 . We will not do so, however. We obtain easier expressions for φ_1 and φ_2 in the following way. We let the derivative e_r act on equation (D.2), then we find using equation (D.2) the relation

$$\varphi_1 + \varphi_2 = \frac{1}{4}(\kappa - \varepsilon)(\mu + \varepsilon). \quad (\text{D.4})$$

At this point, we make a distinction between two cases, $\kappa = \varepsilon$ and $\kappa \neq \varepsilon$. If $\kappa = \varepsilon$, we have $\varphi_1 + \varphi_2 = 0$ and the discussion of case BC in appendix C applies. There, it was shown that this leads to the five-dimensional black hole or a Kasner space. From now on, we assume that $\kappa \neq \varepsilon$. From (D.2) and (D.4), we obtain

$$\varphi_1 = -\frac{1}{8}(\mu + \varepsilon)(\kappa + \mu) \quad \text{and} \quad \varphi_2 = \frac{1}{8}(\mu + \varepsilon)(3\kappa - 2\varepsilon + \mu). \quad (\text{D.5})$$

Because $\varphi_1 \neq 0$, we have $\mu + \varepsilon \neq 0$ and $\kappa + \mu \neq 0$. Inserting (D.5) into the Bianchi identities (B.6), we find

$$e_\alpha(\varepsilon) = -\varepsilon\nu - \mu\nu \quad (\text{D.6}) \quad \text{and} \quad \lambda(\varepsilon - \kappa) + \nu(\varepsilon + \mu) = 0. \quad (\text{D.7})$$

Inserting (D.5) into (B.3), we find

$$(\varepsilon - \kappa)^2 - (\mu - \kappa)^2 + 4\lambda\nu = 0. \quad (\text{D.8})$$

We let the derivative e_α act on (D.7), then we find using (B.5), (D.6), (D.7) and (D.8) the relation

$$\lambda\nu = 0. \quad (\text{D.9})$$

Finally, because we have assumed that $\mu + \varepsilon \neq 0$ and $\kappa \neq \varepsilon$, we get from (D.7) and (D.9) $\nu = 0$ and $\lambda = 0$. From (D.8), (B.5c) and (B.3), we then find $\varepsilon = \mu$. At this point, we have squeezed the differential equations sufficiently. From (B.4) and (B.5) we obtain

$$e_r(\mu) = 2\kappa\mu, \quad e_\alpha(\mu) = 0, \quad e_r(\kappa) = \kappa^2 + \mu^2 \quad \text{and} \quad e_\alpha(\kappa) = 0. \quad (\text{D.10})$$

Because $\delta = 0$, it follows from (B.1d) that we can choose coordinates such that $e_r = \partial_r$ and $e_\alpha = B\partial_\alpha$. The solution of (D.10) is

$$\mu = \frac{1}{2} \left(\frac{1}{r-C} - \frac{1}{r} \right) \quad \text{and} \quad \kappa = -\frac{1}{2} \left(\frac{1}{r-C} + \frac{1}{r} \right),$$

where C is an integration constant. The tetrad e_α can now be determined from (B.1d)

$$e_\alpha = \left(1 - \frac{C}{r} \right)^{1/2} \partial_\alpha.$$

The remaining tetrad vectors can be found using (B.1d) and (B.2)

$$e_t = \left(1 - \frac{C}{r} \right)^{-1/2} \partial_t \quad \text{and} \quad \sigma = (r^2 - Cr)^{-1/2}.$$

This is the metric (4.6).

E. Type 22: solutions in case D, $\varphi_2^2 = \varphi_3^2 + \varphi_4^2$

We now solve the equations (3.1) subject to the condition $\varphi_2^2 = \varphi_3^2 + \varphi_4^2$. We choose our tetrad such that $\varphi_4 = 0$ and $\varphi_3 = -\varphi_2$. From the Bianchi identities (B.6), we obtain two algebraic equations

$$\kappa\varphi_1 + (-2\varepsilon + 3\kappa)\varphi_2 = 0 \quad (\text{E.1}) \quad \text{and} \quad (\delta - \nu)\varphi_2 = 0. \quad (\text{E.2})$$

We will assume that $\varphi_2 \neq 0$, otherwise the discussion in case AD of appendix C shows that we have flat space. Hence, we have $\delta = \nu$. Other equations we obtain from the Bianchi identities and, using (B.3), from equations (B.4) read

$$\begin{aligned} e_r(\varphi_1) &= (5\varepsilon - 9\kappa)\varphi_2 & e_r(\varepsilon) &= \varepsilon^2 + \varepsilon\mu \\ e_r(\varphi_2) &= (\varepsilon + 3\kappa)\varphi_2 & e_r(\kappa) &= \kappa^2 + \kappa\mu \end{aligned} \quad (\text{E.3})$$

We now let the derivative e_r act on equation (E.1). By applying equations (E.3), we find the important relation $\varepsilon(\varepsilon - \kappa)\varphi_2 = 0$. Combining this equation with (E.1), it follows that we have to consider three cases:

- D.1. $\varepsilon = \kappa \neq 0$ and $\varphi_1 + \varphi_2 = 0$,
- D.2. $\varepsilon = 0$, $\kappa \neq 0$ and $\varphi_1 + 3\varphi_2 = 0$,
- D.3. $\varepsilon = \kappa = 0$.

Case D.1: $\varepsilon = \kappa \neq 0$ and $\varphi_1 + \varphi_2 = 0$

This case is treated in case BC of appendix C.

Case D.2: $\varepsilon = 0$, $\kappa \neq 0$ and $\varphi_1 + 3\varphi_2 = 0$

Inserting the relation $\varphi_1 + 3\varphi_2 = 0$ into the Bianchi equations gives $\lambda = -\nu$. From equation (B.3), we get $\varphi_2 = 1/4(\kappa\mu - \nu^2)$. Inserting these results into equations (B.4) and (B.5) gives

$$e_r(\mu) = -3\nu^2 - \mu^2 + 2\kappa\mu \quad (\text{E.4a}) \quad e_\alpha(\mu) = -\nu\mu \quad (\text{E.5a})$$

$$e_r(\nu) = 0 \quad (\text{E.4b}) \quad e_\alpha(\nu) = -\nu^2 \quad (\text{E.5b})$$

$$e_r(\kappa) = \kappa^2 + \kappa\mu \quad (\text{E.4c}) \quad e_\alpha(\kappa) = -\nu\kappa \quad (\text{E.5c})$$

We choose coordinates in such a way that $e_r = A\partial_r$ and $e_\alpha = B\partial_\alpha$. From the commutation relation (B.1d), it follows that B depends only on the coordinate α , hence, by a coordinate transformation, we can put $B = 1$. The general solution of (E.4b) and (E.5b) is $\nu = 1/(\alpha + C_1)$, where C_1 is an arbitrary constant. At this point, it is easier to make a distinction between two cases: $C_1 \neq \infty$ and $C_1 = \infty$.

- $C_1 \neq \infty$

In this case, we can do a coordinate transformation on α to put $C_1 = 0$. The general solution of (E.5) then reads

$$\mu = \frac{f(r)}{\alpha} \quad \text{and} \quad \kappa = \frac{g(r)}{\alpha}.$$

The commutation relation (B.1d) tells us that we can put $A = -\frac{rg(r)}{\alpha}$ by a coordinate transformation on r . The equations (E.4) are now easily solved to yield

$$\kappa = -\frac{1}{\alpha} \left(-1 + \frac{C_3}{r^2} + \frac{2C_2}{r^3} \right)^{1/2} \quad \text{and} \quad \mu = -\frac{r^3 + C_2}{r^3 \alpha} \left(-1 + \frac{C_3}{r^2} + \frac{2C_2}{r^3} \right)^{-1/2}.$$

Finally, the remaining tetrad vectors e_t and e_θ are determined from the commutation relation (B.1a) and the algebraic equation (B.2) respectively

$$e_t = \frac{1}{\alpha r} \left(-1 + \frac{C_3}{r^2} + \frac{2C_2}{r^3} \right)^{-1/2} \partial_t \quad \text{and} \quad \sigma = \frac{\sqrt{C_3}}{\alpha r}.$$

This the metric (4.2).

- $C_1 = \infty$

In this case, $\nu = 0$. The solution of (E.4) and (E.5) can now be found analogously. It reads

$$\begin{aligned} e_r &= \left(C_3 + \frac{2C_2}{r} \right)^{1/2} \partial_r & e_\alpha &= \partial_\alpha \\ \kappa &= -\frac{1}{r} \left(C_3 + \frac{2C_2}{r} \right)^{1/2} & \mu &= -\frac{C_2}{r^2} \left(C_3 + \frac{2C_2}{r} \right)^{-1/2} \end{aligned}$$

The remaining tetrad vectors e_t and e_θ are determined from the commutation relation (B.1a) and the algebraic equation (B.2) respectively

$$e_t = \left(C_3 + \frac{2C_2}{r} \right)^{-1/2} \partial_t \quad \text{and} \quad \sigma = \frac{\sqrt{C_3}}{r}.$$

This is the wrapped black string (2.6).

Case D.3: $\varepsilon = \kappa = 0$

From (B.3), we get $\varphi_1 = \varphi_2 - \lambda\nu$. From (B.4) and (B.5) we obtain the following equations

$$e_r(\mu) = -\nu^2 - \mu^2 + \lambda\nu + 4\varphi_2 \quad (\text{E.6a}) \quad e_\alpha(\mu) = -\nu\mu \quad (\text{E.7a})$$

$$e_r(\nu) = 0 \quad (\text{E.6b}) \quad e_\alpha(\nu) = -\nu^2 + \lambda\nu - 4\varphi_2 \quad (\text{E.7b})$$

$$e_r(\lambda) = 0 \quad (\text{E.6c}) \quad e_\alpha(\lambda) = \lambda^2 + \lambda\nu - 4\varphi_2 \quad (\text{E.7c})$$

If $\nu = 0$, we get from (E.7b) $\varphi_2 = 0$, then $\varphi_1 = 0$ and we obtain flat space. Therefore, we assume in the following that $\nu \neq 0$. We choose coordinates in such a way that $e_r = A\partial_r$ and $e_\alpha = B\partial_\alpha$. From the commutation relation (B.1d), it follows that we can do a coordinate transformation to make A and B both depend only on α . Integrating the equations (E.6a) and (E.7a), we find

$$\mu = \sqrt{C_1} \cot(\sqrt{C_1}r)A \quad \text{and} \quad 4\varphi_2 = \nu^2 - \lambda\nu - C_1A^2,$$

where C_1 is an arbitrary integration constant. We now make a distinction between the two cases $C_1 = 0$ and $C_1 \neq 0$.

- $C_1 = 0$

With hindsight, by a coordinate transformation on α , we put $A = \sinh^{-\frac{1}{\sqrt{3}}} \alpha$. From (B.1d), we obtain $B = \sqrt{3} \nu \tanh \alpha$. The solution of (E.7b) and (E.7c) is now easily found

$$\begin{aligned}\nu &= C_2 \cosh^4 \alpha \sinh^{-2+\frac{2}{\sqrt{3}}} \alpha \\ \lambda &= \frac{1}{\sqrt{3}} \left(6 \tanh^2 \alpha + (-3 + 2\sqrt{3}) \right) \nu\end{aligned}$$

Here, we have used some freedom left over in the choice of coordinates to eliminate one of the two integration constants. The remaining tetrad vectors can be found from (B.2) and (B.1a). All in all, we have

$$\begin{aligned}e_t &= \frac{1}{r} \sinh^{-\frac{1}{\sqrt{3}}} \alpha \partial_t & e_\alpha &= C_2 \sqrt{3} \cosh^3 \alpha \sinh^{-1+\frac{2}{\sqrt{3}}} \alpha \partial_\alpha \\ e_r &= \sinh^{-\frac{1}{\sqrt{3}}} \alpha \partial_r & \sigma &= i C_2 2\sqrt{3} \cosh^2 \alpha \sinh^{-1+\frac{2}{\sqrt{3}}} \alpha\end{aligned}$$

One can see by changing some notation, that this solution is the metric (4.5).

- $C_1 \neq 0$

From (B.2) and (B.1), it follows that the metric can be written as

$$ds^2 = \frac{1}{A(\alpha)^2} \left(dr^2 + \sin^2(\sqrt{C_1} r) dt^2 \right) + \frac{1}{B(\alpha)^2} d\alpha^2 + \frac{1}{\sigma(\alpha)^2} d\Omega^2.$$

This metric belongs to the class (4.3) if $C_1 > 0$, and to the class (4.4) if $C_1 < 0$. We have not been able to find an analytical form of the general solution of (E.6) and (E.7).

References

- [1] R. P. Kerr, “Gravitational Field Of A Spinning Mass As An Example Of Algebraically Special Metrics,” *Phys. Rev. Lett.* **11** (1963) 237.
- [2] J. Stewart, “Advanced General Relativity,” Cambridge University Press, 1994.
- [3] R. C. Myers, “Higher Dimensional Black Holes In Compactified Space-Times,” *Phys. Rev. D* **35** (1987) 455.
- [4] W. Kinnersley, “Type D Vacuum Metrics” *J. Math. Phys.* **10** (1969) 1195.
- [5] M. Carmeli, “Group Theory and General Relativity,” McGraw-Hill Inc., 1977.
- [6] A. Van Proeyen, “Tools for supersymmetry,” arXiv:hep-th/9910030.
- [7] L. Randall and R. Sundrum, “A large mass hierarchy from a small extra dimension,” *Phys. Rev. Lett.* **83** (1999) 3370 [arXiv:hep-ph/9905221]. “An alternative to compactification,” *Phys. Rev. Lett.* **83** (1999) 4690 [arXiv:hep-th/9906064].
- [8] R. Gregory and R. Laflamme, “Black strings and p-branes are unstable,” *Phys. Rev. Lett.* **70** (1993) 2837 [arXiv:hep-th/9301052].

- [9] G. T. Horowitz and K. Maeda, “Fate of the black string instability,” *Phys. Rev. Lett.* **87** (2001) 131301 [arXiv:hep-th/0105111].
- [10] G. T. Horowitz, “Playing with black strings,” arXiv:hep-th/0205069.
- [11] S. S. Gubser, “On non-uniform black branes,” arXiv:hep-th/0110193.
- [12] T. Harmark and N. Obers, “Black holes on cylinders,” arXiv:hep-th/0204047.
- [13] G. W. Gibbons, D. Ida and T. Shiromizu, “Uniqueness and non-uniqueness of static vacuum black holes in higher dimensions,” arXiv:gr-qc/0203004.
- [14] R. Emparan and H. S. Reall, “A rotating black ring in five dimensions,” *Phys. Rev. Lett.* **88** (2002) 101101 [arXiv:hep-th/0110260].
- [15] R. Emparan, G. T. Horowitz and R. C. Myers, “Exact description of black holes on branes,” *JHEP* **0001** (2000) 007 [arXiv:hep-th/9911043].
- [16] A. Chamblin, S. W. Hawking and H. S. Reall, “Brane-world black holes,” *Phys. Rev. D* **61** (2000) 065007 [arXiv:hep-th/9909205].
- [17] R. Emparan and H. S. Reall, “Generalized Weyl solutions,” *Phys. Rev. D* **65** (2002) 084025 [arXiv:hep-th/0110258].
- [18] A. Chodos and S. Detweiler, “Spherically Symmetric Solutions In Five-Dimensional General Relativity,” *Gen. Rel. Grav.* **14** (1982) 879.